

WELCOME TO



OPENDOORS®

Individualized care and support for patients like you™



To start enrollment, text **“START”** to **84537***
or call **1-866-OPENDOOR** (1-866-673-6366)

*Standard data rates may apply.

Visit ofev.com

Please see Important Safety Information throughout this brochure and accompanying Patient Information.



OFEV®
(nintedanib)
capsules 150mg



YOU CAN COUNT ON YOUR



Personal OFEV
CARE TEAM

TO GUIDE YOU THROUGH
your treatment with OFEV® (nintedanib) capsules

As soon as you enroll in OPEN DOORS®, a complimentary program offering compassionate, 24/7 support for you and your caregiver, you'll appreciate all your team can do for you.



To start enrollment, text **"START"** to **84537***
or call **1-866-OPENDOOR (1-866-673-6366)**

*Standard data rates may apply.

What is OFEV?

OFEV is a prescription medicine used to treat people with a lung disease called idiopathic pulmonary fibrosis (IPF). It is not known if OFEV is safe and effective in children.

Important Safety Information

What is the most important information I should know about OFEV (nintedanib)?

OFEV can cause harm, birth defects or death to an unborn baby. Women should not become pregnant while taking OFEV. Women who are able to become pregnant should have a pregnancy test before starting treatment and should use birth control during and for at least 3 months after your last dose. If you become pregnant while taking OFEV, tell your doctor right away.

Please see Important Safety Information throughout this brochure and accompanying [Patient Information](#).



Important Safety Information

What should I tell my doctor before using OFEV?

Before you take OFEV, tell your doctor if you have:

- liver problems
- heart problems
- a history of blood clots
- a bleeding problem or a family history of a bleeding problem
- had recent surgery in your stomach (abdominal) area
- any other medical conditions.

Tell your doctor if you:

- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if OFEV passes into your breast milk. You **should not** breastfeed while taking OFEV.
- are a smoker. You should stop smoking prior to taking OFEV and avoid smoking during treatment.

Please see Important Safety Information throughout this brochure and accompanying [Patient Information](#).

Surrounding you with support



Your benefits include the ability to call or connect with your personal care team virtually anytime. They have a deep understanding of IPF and OFEV® (nintedanib) capsules, and more importantly, they know YOU.

Introducing your personal care team members:



Your Nurse Counselor:

- Helps you understand IPF
- Answers questions about your treatment with OFEV
- Is available 24/7



Your Social Resource Specialist:

- Calls periodically to see how you're doing
- Provides emotional support whenever you need it
- Finds local support for you and/or your caregiver
- Helps to identify local social services, like meal delivery, in-home support, or rides to doctor appointments



Your Case Manager:

- Helps you explore financial assistance options
- Assists in identifying benefits for which you may be eligible
- Coordinates paperwork, like prior authorizations

Important Safety Information

What should I tell my doctor before using OFEV? (cont'd)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements such as St. John's wort.

Please see Important Safety Information throughout this brochure and accompanying [Patient Information](#).

Once you have enrolled in OPEN DOORS[®], your personal care team can connect you with a Clinical Educator.

Your Clinical Educator:

- Meets with you and your family for a personalized (one-on-one or group) educational session
- Helps you understand IPF, how it progresses, and how it will affect you and your family
- Discusses how OFEV[®] works and what to expect on treatment
- Empowers you to play a larger role in your health



Important Safety Information

What are the possible side effects of OFEV?

OFEV may cause serious side effects.

TELL YOUR DOCTOR RIGHT AWAY if you are experiencing any side effects, including:

- **Liver problems.** Unexplained symptoms may include yellowing of your skin or the white part of your eyes (jaundice), dark or brown (tea colored) urine, pain on the upper right side of your stomach area (abdomen), bleeding or bruising more easily than normal, feeling tired, or loss of appetite. Your doctor will do blood tests regularly to check how well your liver is working during your treatment with OFEV.

Please see Important Safety Information throughout this brochure and accompanying [Patient Information](#).

It's easy to enroll



To start enrollment in the OPEN DOORS® Patient Support Program **text "START" to 84537**.* Your personal care representative will reach out to you within the next business day (Monday - Friday 8 AM - 8 PM ET).

*Standard data rates may apply.



Call 1-866-OPENDOOR
(1-866-673-6366)

Important Safety Information

What are the possible side effects of OFEV? (cont'd)

TELL YOUR DOCTOR RIGHT AWAY if you are experiencing any side effects, including:

- **Diarrhea, nausea, and vomiting.** Your doctor may recommend that you drink fluids or take medicine to treat these side effects. Tell your doctor if you have these symptoms, if they do not go away, or get worse and if you are taking over-the-counter laxatives, stool softeners, and other medicines or dietary supplements.
- **Heart attack.** Symptoms of a heart problem may include chest pain or pressure, pain in your arms, back, neck or jaw, or shortness of breath.
- **Stroke.** Symptoms of a stroke may include numbness or weakness on 1 side of your body, trouble talking, headache, or dizziness.

Please see Important Safety Information throughout this brochure and accompanying [Patient Information](#).

It's easy to enroll



Additionally, you can enroll in OPEN DOORS® by filling out the enclosed form and returning it via:



Mail

OPEN DOORS®
P.O. Box 5070
Louisville, KY 40255



Fax

1-844-880-OFEV
(6338)

Important Safety Information

What are the possible side effects of OFEV? (cont'd)

TELL YOUR DOCTOR RIGHT AWAY if you are experiencing any side effects, including:

- **Bleeding problems.** OFEV may increase your chances of having bleeding problems. Tell your doctor if you have unusual bleeding, bruising, or wounds that do not heal and/or if you are taking a blood thinner, including prescription blood thinners and over-the-counter aspirin.
- **Tear in your stomach or intestinal wall (perforation).** OFEV may increase your chances of having a tear in your stomach or intestinal wall. Tell your doctor if you have pain or swelling in your stomach area.

The most common side effects of OFEV are diarrhea, nausea, stomach pain, vomiting, liver problems, decreased appetite, headache, weight loss, and high blood pressure.

These are not all the possible side effects of OFEV. For more information, ask your doctor or pharmacist. **You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <https://www.fda.gov/Safety/MedWatch/default.htm> or call 1-800-FDA-1088.**

Please see Important Safety Information throughout this brochure and accompanying [Patient Information](#).

ENROLL TODAY for round-the-clock access to your personal OFEV[®] care team and all they can do for you.



Text **“START”** to 84537*
or call **1-866-OPENDOOR**
(1-866-673-6366)



*Standard data rates may apply.



Visit ofev.com

Please see Important Safety Information throughout this brochure and accompanying Patient Information.



Copyright © 2019, Boehringer Ingelheim Pharmaceuticals, Inc.
All rights reserved. (2/19) PC-US-107964